

# UNCLASSIFIED

AD NUMBER
ADB269109
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Jun 2001. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St., Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 21 Feb 2003

THIS PAGE IS UNCLASSIFIED

AD \_\_\_\_\_

Award Number: DAMD17-99-1-9134

TITLE: p19 ARF-p53 Tumor Suppressor Pathway During Oncogene-  
Induced Apoptosis and Senescence

PRINCIPAL INVESTIGATOR: Jianping Jin, Ph.D.  
Dr. Scott W. Lowe

CONTRACTING ORGANIZATION: Cold Spring Harbor Laboratory  
Cold Spring Harbor, New York 11724

REPORT DATE: June 2001

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government  
agencies only (proprietary information, Jun 01). Other requests  
for this document shall be referred to U.S. Army Medical Research  
and Materiel Command, 504 Scott Street, Fort Detrick, Maryland  
21702-5012.

The views, opinions and/or findings contained in this report are  
those of the author(s) and should not be construed as an official  
Department of the Army position, policy or decision unless so  
designated by other documentation.

20010810 078

## NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

### LIMITED RIGHTS LEGEND

Award Number: DAMD17-99-1-9134  
Organization: Cold Spring Harbor Laboratory

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

Carol B. Christian

July 30, 2001

**REPORT DOCUMENTATION PAGE**Form Approved  
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

<b>1. AGENCY USE ONLY (Leave blank)</b>		<b>2. REPORT DATE</b> June 2001	<b>3. REPORT TYPE AND DATES COVERED</b> Annual Summary (15 May 00 - 15 May 01)	
<b>4. TITLE AND SUBTITLE</b> p19 ARF-p53 Tumor Suppressor Pathway During Oncogene-Induced Apoptosis and Senescence			<b>5. FUNDING NUMBERS</b> DAMD17-99-1-9134	
<b>6. AUTHOR(S)</b> Jianping Jin, Ph.D. Dr. Scott W. Lowe				
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Cold Spring Harbor Laboratory Cold Spring Harbor, New York 11724  E-Mail: <a href="mailto:jin@cshl.org">jin@cshl.org</a>			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			<b>10. SPONSORING / MONITORING AGENCY REPORT NUMBER</b>	
<b>11. SUPPLEMENTARY NOTES</b>				
<b>12a. DISTRIBUTION / AVAILABILITY STATEMENT</b> Distribution authorized to U.S. Government agencies only (proprietary information, Jun 01). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.				<b>12b. DISTRIBUTION CODE</b>
<b>13. ABSTRACT (Maximum 200 Words)</b>  The primary objective of this study is to provide new insights into the role of the p53 and ARF tumor suppressors in cancer development and therapy. During the second year of this project, we found that E1A oncogene can increase mRNA levels of tumor suppressor ARF. The region of E1A responsible for ARF induction is also important for apoptosis function of E1A. We also found that oncogene E1A can reduce apoptosis inhibitor gene, A20, expression, which can inhibit E1A-induced apoptosis once overexpressed. In collaboration with C. Sher, we tried to produce different monoclonal antibodies to p19ARF. Right now, we are doing the intensive screens to determine the antibodies specificity.				
<b>14. SUBJECT TERMS</b> Breast cancer				<b>15. NUMBER OF PAGES</b> 9
				<b>16. PRICE CODE</b>
<b>17. SECURITY CLASSIFICATION OF REPORT</b> Unclassified	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> Unclassified	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> Unclassified	<b>20. LIMITATION OF ABSTRACT</b> Unlimited	

## Table of Contents

Cover.....	
SF 298.....	II
Table of Contents.....	III
Introduction.....	1
Body.....	2
Key Research Accomplishments.....	4
Conclusions.....	4
References.....	5
Appendices.....	5

## Introduction

Cancer is characterized by dramatic changes in the abundance of proteins that promote growth and metastasis. One of the major changes in tumorigenesis is the loss of the function of tumor suppressor genes. ARF-p53 tumor surveillance pathway has been shown to play important roles in tumor suppression. Mutations in the p53 and other tumor suppressor genes are the most common genetic changes found in breast cancer so far. We and others found that oncogenes induce p53 through a signal transduction pathway that requires the presence of another tumor suppressor, ARF. This pathway, when activated, directs the cells to apoptosis or cell cycle arrest which then prevent tumor cell development. The primary objective of this project is to provide new insights into the action of the tumor suppressor p53 and ARF and the regulation of these two tumor suppressors in cancer development. Our original studies took advantage of mouse cells with a defined genetic background to analyze the relevance of single genes in the oncogene-p19<sup>ARF</sup>-p53 pathway. We proposed to compare this highly controlled system to normal human breast cells in order to determine if the pathway is conserved in human cells and thus potentially relevant to breast cancer. Our early study also indicated that oncogene E1A can sensitize the apoptosis of primary cells which is directly relative to ARF/p53 pathway. Furthermore, by comparing and contrasting normal versus oncogene expressing cells, in these systems, we will be able to get insight to the mechanisms by which p19<sup>ARF</sup> is involved in p53 activation. Finally, a better understanding of the oncogene-p19<sup>ARF</sup>-p53-apoptosis pathway could allow us to exploit new strategies to enhance the chemo- and radio-sensitivity of breast cancer cells. The research progress of the second year on this project is reported as follows.

## Body

### 1. Oncogene signaling to p53 (Aims 2 and 3)

The primary objective of this study is to provide new insights into the role of the p53 and ARF tumor suppressors in breast cancer development and therapy. p53 promotes cell cycle arrest or apoptosis in response to different stimuli, including DNA damage and

mitogenic stimulation. Following DNA damage, signaling is mediated, at least in part, by a kinase that phosphorylates p53 on Serine 15, thereby stabilizing the protein by preventing its association with its negative regulator Mdm2.

In our original proposal, we presented data showing that the adenovirus E1A oncogene activates p53 through a mechanism distinct from DNA damage, involving cancellation of Rb function and activation of the ARF tumor suppressor, leading the cells to apoptosis. ARF-null MEFs expressing E1A are incapable of activating the p53 response and are more resistant to apoptosis following serum depletion or adriamycin treatment, compared to wild type MEFs. Reintroduction of ARF restores p53 accumulation and sensitizes cells to radiation and chemotherapy.

During the second year of this research, we decided to test whether ARF induction is a common event following oncogene overexpression. We found that E1A can induce ARF expression both in human and mouse primary cells. More importantly, northern blot indicated that E1A-induced ARF expression results from the increase of ARF mRNA levels.

Our published and unpublished data indicated that different regions of E1A correspond to various function of E1A. For example, the CR2 region has been shown to bind to RB and be essential to apoptosis function of E1A (Samuelson et al., 1997). We also found that p300 binding region which locates at N-terminal part of E1A is not important to the apoptosis function of E1A. However, region 26-35 is important. Now, we are focusing on which regions of E1A polypeptide are responsible for the increase of ARF mRNA levels by E1A in primary cells. We expected that the regions important for E1A's apoptosis function should be also important for ARF induction. Our preliminary data suggested that both CR2 and 26-35 regions are important for the increase of ARF mRNA levels. In collaboration with Dr. David Livingston's Lab at Harvard University, we found that the 26-35 region binds p400/TRAPP complex which possibly involved in ARF induction at transcription levels (Samuelson et al., unpublished). Next, we will focus on 26-35 region for ARF induction. Our data indicated that region 26-35 does not completely lose the function of E1A in apoptosis induction. Therefore, we will make more mutants around this region to study ARF induction more deeply.

## 2. Production of highly specific monoclonal antibodies against the N-terminal portion of ARF(Aim 1)

In human cancers, ARF mutations occur mainly in the region coding for the C-terminus domain (exon 2), possibly leading to the formation of truncated isoforms. These putative forms would be impossible to be detected by the currently available anti-ARF antibodies, which are directed against the C-terminus of the protein. In the original proposal, we proposed to raise antibodies against the N-terminal region of ARF, which has been shown to be necessary and sufficient to induce cell cycle arrest and to interact with p53 and/or Mdm2.

In collaboration with C. Sherr, we produced and analyzed different monoclonal antibodies. The CSHL Monoclonal Service Facility performed all the animal work, the fusions, single cell cloning and initial screens. We are currently doing the extensive screens to determine the antibodies specificity. If we found the specific antibodies for ARF, we will initially use them for routine characterization studies, such as western blot analysis and immunofluorescence, to detect ARF levels in different cell lines and to investigate the presence and abundance of the truncated isoforms in tumor derived cell lines and in clinical specimens.

## 3. The function of E1A in suppression of anti-apoptosis proteins

Beside the induction of positive apoptosis effector by oncogenes, I am also interested in the regulation of apoptosis inhibitors by oncogenes. We found that E1A can repress A20, an apoptosis inhibitor, using a Clontech Atlas filter array. Since A20 has been shown to be overexpressed in undifferentiated nasopharyngeal carcinoma, head & neck squamous cell carcinomas, it would be interesting to see whether A20 could be one of the reasons that many cancers are resistant to cancer drug treatments and whether A20 would be a good drug target for the cancers with high A20 expression. I confirmed that repression of A20 by E1A exists generally in human and mouse primary cells by northern blot. Interestingly, it is possible that E1A represses A20 expression by affecting A20 mRNA stability. Now, I have started to determine whether other oncogenes could have similar function as E1A in inhibition of A20. I also made A20/retrovirus by retrovirus system, I infected the mammalian primary cells with these A20/retrovirus, I was able to



show that A20 overexpression could block oncogene/cancer drug-induced apoptosis in some cells.

### **Key Research Accomplishments**

We found that E1A can induce ARF expression by transcription and/or post-transcription regulation. Consequently, The function of Mdm2 is inhibited and tumor suppress p53 be stabilized by elevated ARF protein levels. More importantly, this induction of ARF by E1A is correlated to apoptosis induction by E1A in primary cells. The region from residue 26 to 35 and conserved region 2 (CR2) of E1A 12S polypeptide is important to apoptosis function of E1A. Our data indicated that these two regions are responsible for ARF induction by E1A at mRNA levels.

We also examined the role of oncogene E1A in regulation of an apoptosis inhibitor, A20. We found oncogene E1A can inhibit A20 expression at mRNA level. Most interesting, overexpression of A20 can block E1A/cancer drug-induced apoptosis.

In collaboration with C. Sherr, we tried to produce and analyze different monoclonal antibodies to p19<sup>ARF</sup>. Right now, we are in the process of intensive screens to determine the antibodies specificity.

### **Reportable Outcomes**

The research is still ongoing. In collaboration with Mr. Samuelson, we will have a paper about E1A-induced ARF expression in a short time. The paper is in writing. At this moment, there is no reportable outcome.

### **Conclusions**

In conclusion, oncogenes can stabilize p53 level by activating ARF at mRNA levels. In collaboration with Mr. Samuelson, I found that oncogene E1A induces ARF mRNA

expression. Mutational analysis indicated that the regions of E1A polypeptide responsible for ARF induction are also important to apoptosis function of E1A. These data supplied the solid support for ARF-p53 pathway in tumor surveillance. How E1A induce ARF expression is under investigation in our Laboratory. In collaboration with Dr. de Stanchina, I also E1A can repress the expression of an apoptosis inhibitor, A20, beside activate ARF-p53 pathway. Our preliminary data indicated that the repression of A20 by oncogene E1A could be a general phenomena. Oncogenes, E2F and myc may have similar function. We are continuing to work on this project, with a particular emphasis on how E1A induces ARF expression.

### **References**

N/A

### **Appendices**

N/A



DEPARTMENT OF THE ARMY  
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND  
504 SCOTT STREET  
FORT DETRICK, MARYLAND 21702-5012

REPLY TO  
ATTENTION OF:

MCMR-RMI-S (70-1y)

21 Feb 03

MEMORANDUM FOR Administrator, Defense Technical Information  
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,  
VA 22060-6218


SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for this Command. Request the limited distribution statement for the enclosed accession numbers be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Kristin Morrow at DSN 343-7327 or by e-mail at Kristin.Morrow@det.amedd.army.mil.

FOR THE COMMANDER:

Encl

  
PHYLLIS M. RINEHART  
Deputy Chief of Staff for  
Information Management

ADB263458	ADB282838
ADB282174	ADB233092
ADB270704	ADB263929
ADB282196	ADB282182
ADB264903	ADB257136
ADB268484	ADB282227
ADB282253	ADB282177
ADB282115	ADB263548
ADB263413	ADB246535
ADB269109	ADB282826
ADB282106	ADB282127
ADB262514	ADB271165
ADB282264	ADB282112
ADB256789	ADB255775
ADB251569	ADB265599
ADB258878	ADB282098
ADB282275	ADB232738
ADB270822	ADB243196
ADB282207	ADB257445
ADB257105	ADB267547
ADB281673	ADB277556
ADB254429	ADB239320
ADB282110	ADB253648
ADB262549	ADB282171
ADB268358	ADB233883
ADB257359	ADB257696
ADB265810	ADB232089
ADB282111	ADB240398
ADB273020	ADB261087
ADB282185	ADB249593
ADB266340	ADB264542
ADB262490	ADB282216
ADB266385	ADB261617
ADB282181	ADB269116
ADB262451	
ADB266306	
ADB260298	
ADB269253	
ADB282119	
ADB261755	
ADB257398	
ADB267683	
ADB282231	
ADB234475	
ADB247704	
ADB258112	
ADB267627	